


BASIC MATHEMATICS



Trigonometry

• **Angle** : The angle covered by the revolving line OP is $\theta = \angle POX$



$1^\circ = 60'$ (minute) ; $1' = 60''$ (second)

$\Rightarrow 1 \text{ rad} = \frac{180^\circ}{\pi} \approx 57.3^\circ$

\Rightarrow Angle θ° to Radian multiply it by $\frac{\pi}{180^\circ}$.

\Rightarrow Angle Radian to θ° multiplying it by $\frac{180^\circ}{\pi}$.


• **Trigonometrical Ratios** :

$\sin \theta = \frac{P}{H} = \frac{MP}{OP}$ $\cos \theta = \frac{B}{H} = \frac{OM}{OP}$

$\tan \theta = \frac{P}{B} = \frac{MP}{OM}$ $\cot \theta = \frac{B}{P} = \frac{OM}{MP}$

$\sec \theta = \frac{H}{B} = \frac{OP}{OM}$ $\text{cosec } \theta = \frac{H}{P} = \frac{OP}{MP}$

$\Rightarrow \sin^2 \theta + \cos^2 \theta = 1$; $1 + \tan^2 \theta = \sec^2$; $1 + \cot^2 \theta = \text{cosec}^2 \theta$



• **Table : Trigonometry Standard angles from 0° to 180°**

• **Four Quadrants and ASTC Rule** :

In Ist quadrant, all trigonometric ratios are positive.

\Rightarrow In IInd quadrant, only $\sin \theta$ and $\text{cosec } \theta$ are positive.

\Rightarrow In IIIrd quadrant, only $\tan \theta$ and $\cot \theta$ are positive.

\Rightarrow In IVth quadrant, only $\cos \theta$ and $\sec \theta$ are positive.

| | | | |
|----------------------------|------|---------------------------|-----|
| | 90° | | |
| II nd quadrant | sin | I st quadrant | All |
| 180° | | | 0° |
| | | tan | cos |
| III rd quadrant | | IV th quadrant | |
| | 270° | | |

• **Important trigonometric formula** :

| Angle (θ) | 0 | 30° | 45° | 60° | 90° | 120° | 135° | 150° | 180° |
|--------------------|---|----------------------|----------------------|----------------------|---------------|----------------------|-----------------------|-----------------------|------|
| $\sin \theta$ | 0 | $\frac{1}{2}$ | $\frac{1}{\sqrt{2}}$ | $\frac{\sqrt{3}}{2}$ | 1 | $\frac{\sqrt{3}}{2}$ | $\frac{1}{\sqrt{2}}$ | $\frac{1}{2}$ | 0 |
| $\cos \theta$ | 1 | $\frac{\sqrt{3}}{2}$ | $\frac{1}{\sqrt{2}}$ | $\frac{1}{2}$ | 0 | $-\frac{1}{2}$ | $-\frac{1}{\sqrt{2}}$ | $-\frac{\sqrt{3}}{2}$ | -1 |
| $\tan \theta$ | 0 | $\frac{1}{\sqrt{3}}$ | 1 | $\sqrt{3}$ | (not defined) | $-\sqrt{3}$ | -1 | $-\frac{1}{\sqrt{3}}$ | 0 |


• **Range of trigonometric functions** :

$\Rightarrow \sin \theta = \frac{P}{H}$ and $P \leq H$

So, $-1 \leq \sin \theta \leq 1$

$\Rightarrow \cos \theta = \frac{B}{H}$ and $B \leq H$, So $-1 \leq \cos \theta \leq 1$


$\Rightarrow \tan \theta = \frac{P}{B}$ So, $-\infty < \tan \theta < \infty$



• **Small Angle Approximation** :

If θ is small ($\theta < 45^\circ$)


$\sin \theta \approx \theta$; $\cos \approx 1$ and $\tan \theta \approx \theta$



Coordinate Geometry

• **Origin** : ANY fixed point from which all measurements are taken from this.

• **Axis** : ANY fixed direction passing through origin.




• **Distance Formula** :

$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

In 3-d (space) - $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$

• **Slope of a line** :

$m = \tan \theta = \frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1}$




• **Important formulae of Differentiation** :

$\Rightarrow \frac{d}{dx}(\sin x) = \cos x \Rightarrow \frac{d}{dx}(\cos x) = -\sin x$ (#) $\frac{d}{dx}(\log_e x) = \frac{1}{x}$

$\Rightarrow \frac{d}{dx}(\tan x) = \sec^2 x \Rightarrow \frac{d}{dx}(\cot x) = -\text{cosec}^2 x$ (#) $\frac{d}{dx}(e^x) = e^x$

$\Rightarrow \frac{d}{dx}(\text{cosec } x) = -\text{cosec } x \cot x$ (#) $\frac{d}{dx}(e^{ax}) = ae^x$



• **Concept of Maxima and Minima**





\Rightarrow Condition for minima : $\left[\frac{dy}{dx} = 0 \text{ and } \frac{d^2y}{dx^2} > 0 \right]$

\Rightarrow Condition for maxima : $\left[\frac{dy}{dx} = 0 \text{ and } \frac{d^2y}{dx^2} < 0 \right]$

Algebra

• **Quadratic Equation and its Solutions** :
AN algebraic equation of 2nd order is called a quadratic equation. Equation = $ax^2 + bx + c = 0$

General Solution $\Rightarrow x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$



Differentiation

• **Physical Meaning of $\frac{dy}{dx}$**

(i) The ratio of small change in the function y and the variable x is called the average rate of change y w.r.t. x.

(ii) When $\Delta x \rightarrow 0$. The limiting value of $\frac{\Delta y}{\Delta x}$ is $\lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x} = \frac{dy}{dx}$

• **Main Formulas of Differentiation**

1. $\frac{d}{dx}(K) = 0$ K = constant

2. $\frac{d}{dx}(KU) = K \frac{dU}{dx}$ [U is a function of x]


3. $\frac{d}{dx}(U \pm V \pm W) = \frac{dU}{dx} \pm \frac{dV}{dx} \pm \frac{dW}{dx}$

where U, V and W are functions of x.

4. $\frac{d}{dx}(UV) = U \frac{dV}{dx} + V \frac{dU}{dx}$

5. $\frac{d}{dx}\left(\frac{U}{V}\right) = \frac{V \frac{dU}{dx} - U \frac{dV}{dx}}{V^2}$

6. $\frac{d}{dx}(x^n) = nx^{n-1}$



gives $x_1 = \frac{b + \sqrt{b^2 - 4ac}}{2a}$ and $x_2 = \frac{-b - \sqrt{b^2 - 4ac}}{2a}$

$\Rightarrow b^2 - 4ac \geq 0$ For real roots.

$\Rightarrow b^2 - 4ac < 0$ For Imaginary roots.

• **Binomial Expression** :

AN algebraic expression having two terms only. **LOREM IPSUM**

Example : $(a + b)$, $(a + b)^4$, $(2x - 3y)^2$ etc.

• **Binomial Theorem** :

$(a + b)^n = a^n + na^{n-1}b + \frac{n(n-1)}{2}a^{n-2}b^2 + \dots$

Standard Graphs and their Equations

Integration

If I is the integration of $f(x)$ with respect to x then $I = \int f(x) dx$

• **Main Formulae of Integration** :


1. $\int x^n dx = \frac{x^{n+1}}{n+1} + c$, $n \neq -1$

2. $\int \sin x dx = -\cos x + c$, c = constant

3. $\int \cos x dx = \sin x + c$

4. $\int \frac{1}{x} dx = \log_e x + c$

5. $\int e^x dx = e^x + c$



• **Define Integrals** :

If $\frac{d}{dx}(f(x)) = f'(x)$ then $\int_a^b f'(x) dx = [f(x)]_a^b$

is called definite integral.

• **Area Under Curve** :

$\int_a^b f(x) dx =$ Shaded area between curve and x-axis.

$(1 + x)^n = 1 + nx + \frac{n(n-1)x^2}{2 \times 1} + \dots$

• **Logarithm Main Formulae** :

$\log mn = \log m + \log n$ $\log m^n = n \log m$

$\log \frac{m}{n} = \log m - \log n$ $\log_e m = 2.303 \log_{10} m$

• **Arithmetic Progression (AP)** :

(AP) = $a + a + d + a + d + \dots + a + (n-1)d$
where a = first term d = common difference

\Rightarrow Sum of n term's $S_n = \frac{n}{2}[2a + (n-1)d] = \frac{n}{2}[a + n^{\text{th}} \text{ term}]$

(i) Sum of first n natural number's $S_n = \frac{n(n+1)}{2}$

(ii) Sum of squares of first n $S_n = \frac{n(n+1)(2n+1)}{6}$

• **Geometric Progression (GP)**

(GP) = $a, ar, ar^2, ar^3, \dots, ar^{n-1}$; a = first term, r = common ratio

Sum of n terms $\rightarrow S_n = \frac{a(1-r^n)}{1-r}$; For $0 \leq |r| < 1$

Sum of ∞ terms $S_\infty = \frac{a}{1-r}$